

INEQUALITY FOR VOICULESCU'S FREE ENTROPY IN TERMS OF BROWN MEASURE

PIOTR ŚNIADY

ABSTRACT. We study Voiculescu's microstate free entropy for a single non-selfadjoint random variable. The main result is that certain additional constraints on eigenvalues of microstates do not change the free entropy. Our tool is the method of random regularization of Brown measure which was studied recently by Haagerup and the author. As a simple application we present an upper bound for the free entropy of a single non-selfadjoint operator in terms of its Brown measure and the second moment. We furthermore show that this inequality becomes an equality for a class of DT -operators which was introduced recently by Dykema and Haagerup.

1. INTRODUCTION

The microstate free entropy χ was introduced by Voiculescu [Voi94] as a tool for the study of some non-commutative systems. Roughly speaking, it answers the question how many finite matrices have nearly the same moments as a given non-commutative random variable. It has turned out to have very powerful applications (cf. [Ge97, Voi96]), however it is not an easy object to deal with.

One of the reasons of these difficulties is that currently there are no general methods for the computation of the free entropy in concrete cases. Exact formulas were found for the free entropy of a single selfadjoint random variable, for tuples of free random variables [Voi94] and for R -diagonal elements [NSS99].

In this article we present a method which hopefully will be useful for calculating and estimating free entropy in many concrete cases. The main idea is to change the definition of microstates Γ which approximate a single non-selfadjoint random variable x in such a way that it does not change the value of the free entropy $\chi(x)$.

The original sets Γ consisted of all matrices which—informally speaking—had almost the same moments as a given random variable x , while our new sets $\tilde{\Gamma}$ will consist of these matrices in Γ which additionally have similar eigenvalues to the Brown measure of x . In order to show that Γ

and $\tilde{\Gamma}$ give rise to the same free entropy we use the method of random regularization of Brown spectral measure which was introduced by Haagerup [Haa01] and further developed by the author [Śni01].

As an application we present a new upper bound for the free entropy of a single random variable x in terms of its Brown measure and second moment. We show also that for a class of DT -operators which was introduced recently by Dykema and Haagerup [DH01] this inequality becomes equality.

2. PRELIMINARIES

2.1. Non-commutative probability spaces. A non-commutative probability space is a pair $(\mathcal{A}, \phi)$, where $\mathcal{A}$ is a $\star$ -algebra and ϕ is a normal, faithful, tracial state on $\mathcal{A}$. Elements of $\mathcal{A}$ will be referred to as non-commutative random variables and state ϕ as expectation value.

One of the simplest examples is the set $\mathcal{M}_N$ of all complex-valued $N \times N$ matrices equipped with a normalized trace tr given by $\text{tr } m = \frac{1}{N} \text{Tr } m$, where $m \in \mathcal{M}_N$ and Tr is the usual trace.

2.2. Microstate free entropy. The original definition of Voiculescu's free entropy $\chi^{\text{sa}}(x_1, \dots, x_n)$ allowed to compute the free entropy of a tuple of non-commutative self-adjoint random variables. The considered in this article free entropy $\chi(x)$ of a non-selfadjoint random variable is connected with the original definition by

$$\chi(x) = \chi^{\text{sa}}(\Re x, \Im x).$$

Let x be a non-commutative random variable, $\epsilon > 0$, $R > 0$ be real numbers and $k > 0$ be integer. We define the sets [Voi94, NSS99]

$$(1) \quad \Gamma_R(x; k, N, \epsilon) = \left\{ m \in \mathcal{M}_N : \|m\| \leq R \text{ and } \right. \\ \left. |\text{tr}(m^{s_1} \dots m^{s_p}) - \tau(x^{s_1} \dots x^{s_p})| < \epsilon \right. \\ \left. \text{for all } p \leq k \text{ and } s_1, s_2, \dots, s_p \in \{1, \star\} \right\}.$$

Define next

$$(2) \quad \chi_R(x; k, \epsilon) = \limsup_{N \rightarrow \infty} \left[\frac{1}{N^2} \log \text{vol } \Gamma_R(x; k, N, \epsilon) + \log N \right],$$

where vol is a Lebesgue measure on $\mathcal{M}_N$ as described in (10). Lastly, the free entropy is defined by

$$(3) \quad \chi(x) = \sup_R \inf_{k, \epsilon} \chi_R(x; k, \epsilon).$$

Since $\chi_R(x; k, \epsilon)$ is a decreasing function of k and an increasing function of ϵ , hence we have the following simple lemma.

Lemma 1. *Let a non-commutative random variable x and a number $R > 0$ be given. Then there exists a sequence (ϵ_N) of non-negative numbers and a sequence (k_N) of natural numbers such that $\lim_{N \rightarrow \infty} \epsilon_N = 0$, $\lim_{N \rightarrow \infty} k_N = \infty$ and*

$$(4) \quad \chi_R(x) \leq \limsup_{N \rightarrow \infty} \chi_R(x; k_N, N, \epsilon_N).$$

2.3. Fuglede–Kadison determinant and Brown measure. Let a non-commutative probability space $(\mathcal{A}, \phi)$ be given. For $x \in \mathcal{A}$ we define its Fuglede–Kadison determinant $\Delta(x)$ by [FK52]

$$\Delta(x) = \exp[\phi(\ln |x|)]$$

and its Brown measure [Bro86] to be the Schwartz distribution on $\mathbb{C}$ given by

$$\mu_x = \frac{1}{2\pi} \left(\frac{\partial^2}{\partial(\Re \lambda)^2} + \frac{\partial^2}{\partial(\Im \lambda)^2} \right) \ln \Delta(x - \lambda).$$

One can show that in fact μ_x is a positive probability measure on $\mathbb{C}$.

Lemma 2. *The Brown measure of a matrix $m \in \mathcal{M}_N$ with respect to the state tr is a probability counting measure on the set of eigenvalues of m :*

$$\mu_m = \frac{1}{N} \sum_{i=1}^N \delta_{\lambda_i},$$

where $\lambda_1, \dots, \lambda_N$ are the eigenvalues of m counted with multiples.

In the following we will be interested in studying the random measure $\omega \mapsto \mu_{A(\omega)}$ for a random matrix $A \in \mathcal{M}_N(\mathcal{L}^{\infty-}(\Omega))$. This random measure is called the empirical eigenvalues distribution.

2.4. Convergence of $\star$ -moments. Let a sequence (A_N) of random matrices (where $A_N \in \mathcal{M}_N(\mathcal{L}^{\infty-}(\Omega))$), a non-commutative probability space $(\mathcal{A}, \phi)$ and $x \in \mathcal{A}$ be given. We say that the sequence A_N converges to x in $\star$ -moments almost surely if for every $n \in \mathbb{N}$ and $s_1, \dots, s_n \in \{1, \star\}$ we have that

$$\lim_{N \rightarrow \infty} \text{tr}_N[A_N^{s_1} \cdots A_N^{s_n}] = \phi(x^{s_1} \cdots x^{s_n})$$

holds almost surely.

2.5. Random regularization of Brown measure. We say that a random matrix

$$G_N = (G_{N,ij})_{1 \leq i,j \leq N} \in \mathcal{M}_N(\mathcal{L}^{\infty-}(\Omega))$$

is a standard Gaussian random matrix if

$$(\Re G_{N,ij})_{1 \leq i,j \leq N}, (\Im G_{N,ij})_{1 \leq i,j \leq N}$$

are independent Gaussian variables with mean zero and variance $\frac{1}{2N}$.

Theorem 3. Let (A_N) be a sequence of random matrices, $A_N \in \mathcal{M}_N(\mathcal{L}^{\infty-}(\Omega))$, which converges in $\star$ -moments to x almost surely.

Let furthermore (G_N) be a sequence of independent standard Gaussian matrices which is independent of (A_N) . There exists a sequence (t_N) of real numbers such that $\lim_{N \rightarrow \infty} t_N = 0$ and such that the sequence of empirical eigenvalues distributions $\mu_{A_N + t_N G_N}$ converges weakly to μ_x almost surely.

There also exists a sequence (B_N) of non-random matrices $B_N \in \mathcal{M}_N$ such that $\lim_{N \rightarrow \infty} \|B_N\| = 0$ and such that the sequence of empirical eigenvalues distributions $\mu_{A_N + B_N}$ converges weakly to μ_x almost surely.

Proof. The first part was proved in [Śni01].

For the second part of the theorem let us define $B_N = t_N G_N(\omega)$. Since $\limsup_{N \rightarrow \infty} \|G_N\| < \infty$ holds almost surely [Gem80], hence so defined sequence B_N fulfills the hypothesis of the theorem almost surely. $\square$

3. THE MAIN RESULT: IMPROVED MICROSTATES $\tilde{\Gamma}$

Let x be a non-commutative random variable, $\epsilon > 0$, $R > 0$ be real numbers and $k > 0$, $m > 0$ be integers. In the full analogy with (1)–(3) we define improved microstates $\tilde{\Gamma}$ and improved free entropy $\tilde{\chi}$:

$$(5) \quad \tilde{\Gamma}_R(x; k, N, \epsilon, l, \theta) = \left\{ m \in \Gamma_R(x; k, N, \epsilon) : \left| \int_{\mathbb{C}} z^i \bar{z}^j d\mu_m - \int_{\mathbb{C}} z^i \bar{z}^j d\mu_x \right| < \theta \quad \text{for } i, j \leq l \right\},$$

$$(6) \quad \tilde{\chi}_R(x; k, \epsilon, l, \theta) = \limsup_{N \rightarrow \infty} \left[\frac{1}{N^2} \log \text{vol } \tilde{\Gamma}_R(x; k, N, \epsilon, l, \theta) + \log N \right],$$

$$(7) \quad \tilde{\chi}(x) = \sup_R \inf_{k, \epsilon, l, \theta} \tilde{\chi}_R(x; k, \epsilon, l, \theta).$$

Theorem 4. For every non-commutative random variable x we have

$$\chi(x) = \tilde{\chi}(x).$$

Proof. Since $\tilde{\Gamma}_R(x; k, N, \epsilon, l, \theta) \subseteq \Gamma_R(x; k, N, \epsilon)$ hence $\tilde{\chi}(x) \leq \chi(x)$ follows easily.

Let (ϵ_N) and (k_N) be the sequences given by Lemma 1. Let (A_N) be a sequence of independent random matrices such that the distribution of A_N is the uniform distribution on the set $\Gamma_R(x; k_N, N, \epsilon_N)$ and let (B_N) be the sequence given by Theorem 3.

Since $\|B_N\|$ converges to zero, hence there exists $R' > 0$, a sequence of positive numbers (ϵ'_N) which converges to zero and a sequence of natural

numbers (k'_N) which diverges to infinity such that

$$\Gamma_R(x; k_N, N, \epsilon_N) + B_N \subseteq \Gamma_{R'}(x; k'_N, N, \epsilon'_N)$$

holds for every $N \in \mathbb{N}$, where $\Gamma_R(x; k_N, N, \epsilon_N) + B_N$ denotes a translation of the set $\Gamma_R(x; k_N, N, \epsilon_N)$ by the vector B_N .

Since random measures $\omega \mapsto \mu_{A_N(\omega) + B_N}$ converge weakly to μ_x in probability, hence for any $\theta > 0$ and integer $l > 0$ we have that

$$\lim_{N \rightarrow \infty} P(\omega : A_N(\omega) + B_N \notin \tilde{\Gamma}_{R'}(x; k'_N, N, \epsilon'_N, l, \theta)) = 0.$$

Since the Lebesgue measure is translation-invariant, hence for any $\theta > 0$ and integer $l > 0$ we have

$$\limsup_{N \rightarrow \infty} \frac{\text{vol } \Gamma_R(x; k_N, N, \epsilon_N)}{\text{vol } \tilde{\Gamma}_{R'}(x; k'_N, N, \epsilon'_N, l, \theta)} \leq 1,$$

or equivalently

$$(8) \quad \limsup_{N \rightarrow \infty} (\chi_R(x; k_N, N, \epsilon_N) - \tilde{\chi}_{R'}(x; k'_N, N, \epsilon'_N, l, \theta)) \leq 0.$$

For any $\epsilon > 0$ and integer $k > 0$ there exists N_0 such that for any $N > N_0$ we have $\epsilon_N < \epsilon$ and $k_N > k$, hence for $N > N_0$

$$(9) \quad \tilde{\chi}_{R'}(x; k'_N, N, \epsilon'_N, l, \theta) \leq \tilde{\chi}_{R'}(x; k, N, \epsilon, l, \theta).$$

Inequalities (4), (8) and (9) combine to give

$$\chi(x) \leq \tilde{\chi}(x).$$

□

4. APPLICATION: UPPER BOUND FOR FREE ENTROPY

In this section we present a new inequality for the free entropy of a single non-selfadjoint random variable. The main idea is to write matrices from microstates $\tilde{\Gamma}$ in the upper triangular form and then to find constraints on diagonal and offdiagonal entries.

4.1. Pull-back of the Lebesgue measure on $\mathcal{M}_N$. We denote by $\mathcal{M}_N^d = \{m \in \mathcal{M}_N : m_{ij} = 0 \text{ if } i \neq j\}$ the set of diagonal matrices and by $\mathcal{M}_N^{\text{sut}} = \{m \in \mathcal{M}_N : m_{ij} = 0 \text{ if } i \geq j\}$ the set of all strictly upper triangular matrices.

We can regard $\mathcal{M}_N$ and $\mathcal{M}_N^{\text{sut}}$ as real Euclidean spaces with a scalar product $\langle x, y \rangle = \Re \text{Tr } xy^*$ and thus equip them with Lebesgue measures

$$(10) \quad \text{vol} = \prod_{1 \leq i, j \leq N} d\Re m_{ij} d\Im m_{ij}$$

and

$$\text{vol}^{\text{sut}} = \prod_{1 \leq i < j \leq N} d\Re m_{ij} d\Im m_{ij}$$

respectively.

We have a clear isomorphism $\mathcal{M}_N^{\text{d}} = \{(\lambda_1, \dots, \lambda_N) \in \mathbb{C}^N\}$ and we equip it with a measure

$$\text{vol}^{\text{d}} = \frac{\pi^{\frac{N^2-N}{2}}}{\prod_{1 \leq i \leq N} i!} \prod_{1 \leq i < j \leq N} |\lambda_i - \lambda_j|^2 \prod_{1 \leq i \leq N} d\Re \lambda_i d\Im \lambda_i.$$

We also denote by U_N the set of unitary $N \times N$ matrices equipped with the Haar measure vol^{U} normalized in such a way that it is a probability measure.

Proposition 5. *For every N the measure $\text{vol}^{\text{d}} \times \text{vol}^{\text{sut}} \times \text{vol}^{\text{U}}$ is a pull-back of the measure vol with respect to the map*

$$\mathcal{M}_N^{\text{d}} \times \mathcal{M}_N^{\text{sut}} \times U_N \ni (d, m, u) \mapsto u(d + m)u^{-1} \in \mathcal{M}_N.$$

Proof. This result is due to Dyson and can be extracted from Appendix A.35 of [Meh91]. $\square$

4.2. Diagonal entropy $\hat{\chi}^{\text{d}}$. Let x be a non-commutative random variable. In the following we define an auxiliary quantity $\hat{\chi}^{\text{d}}(\nu)$ which would answer the question how many diagonal matrices (with respect to the measure vol^{d}) have almost the same Brown measure as x .

$$\begin{aligned} \hat{\Gamma}_R^{\text{d}}(x; N, l, \theta) &= \left\{ m \in \mathcal{M}_N^{\text{d}} : \|m\| \leq R \text{ and } \right. \\ &\quad \left. \left| \int_{\mathbb{C}} z^i \bar{z}^j d\mu_m - \int_{\mathbb{C}} z^i \bar{z}^j d\mu_x \right| < \theta \text{ for all } i, j \leq l \right\}; \\ \hat{\chi}_R^{\text{d}}(x; l, \theta) &= \lim_{N \rightarrow \infty} \left[\frac{1}{N^2} \log \text{vol}^{\text{d}} \hat{\Gamma}_R^{\text{d}}(x; l, \theta) + \frac{\log N}{2} \right], \\ \hat{\chi}^{\text{d}}(x) &= \sup_R \inf_{l, \theta} \hat{\chi}_R^{\text{d}}(x; l, \theta). \end{aligned}$$

Theorem 6. *For any non-commutative random variable x we have*

$$\hat{\chi}^{\text{d}}(x) = \int_{\mathbb{C}} \int_{\mathbb{C}} \log |z_1 - z_2| d\mu_x(z_1) d\mu_x(z_2) + \frac{3}{4} + \frac{\ln \pi}{2}.$$

Proof. Proof follows exactly the proof of Proposition 4.5 of [Voi94], but since we are dealing with measures on $\mathbb{C}$ and the original proof concerns measures on $\mathbb{R}$, we have to replace Lemma 4.3 in [Voi94] by Theorem 2.1 of [Had98]. $\square$

4.3. Offdiagonality. If x is a non-commutative random variable we define its offdiagonality od_x by

$$\text{od}_x = \tau(xx^*) - \int_{\mathbb{C}} |z|^2 d\mu_x(z).$$

This quantity can be regarded as a kind of a non-commutative variance. Since $\text{od}_x = 0$ if and only if x is normal, hence offdiagonality of x can be also regarded as a kind of a distance of x to normal operators.

For an upper-triangular matrix $m \in \mathcal{M}_N(\mathbb{C})$ ($m_{ij} = 0$ if $i > j$) its offdiagonality is equal to the (normalized) sum of squares of the offdiagonal entries:

$$\text{od}_m = \frac{1}{N} \sum_{1 \leq i < j \leq N} |m_{ij}|^2.$$

Proposition 7. *For any $o > 0$ we have that*

$$\lim_{N \rightarrow \infty} \left[\frac{1}{N^2} \log \text{vol}^{\text{sut}} \{m \in \mathcal{M}_N^{\text{sut}} : \text{od}_m \leq o\} + \frac{\log N}{2} \right] = \frac{1}{2} + \frac{\log 2\pi o}{2}.$$

Proof. It is enough to notice that $\{m \in \mathcal{M}_N^{\text{sut}} : \text{tr } mm^* \leq o\}$ is a $N(N-1)$ -dimensional ball with radius $\sqrt{oN}$, hence its volume is equal to

$$\pi^{\frac{N(N-1)}{2}} \left[\Gamma \left(\frac{N(N-1)}{2} + 1 \right) \right]^{-1} (oN)^{\frac{N(N-1)}{2}}.$$

□

4.4. The main inequality.

Theorem 8. *Let x be a non-commutative random variable. Then*

$$(11) \quad \chi(x) \leq \int_{\mathbb{C}} \int_{\mathbb{C}} \log |z_1 - z_2| d\mu_x(z_1) d\mu_x(z_2) + \frac{5}{4} + \ln \pi \sqrt{2 \text{od}_x}.$$

Proof. Proof is a direct consequence of Theorem 4, Proposition 5, Theorem 6 and Proposition 7. □

4.5. Free entropy of DT -operators. For any compactly supported probability measure ν on $\mathbb{C}$ and $o \geq 0$ Dykema and Haagerup consider an operator x which is said to be $DT(\nu, o)$ [DH01]. This operator is implicitly defined to be the expected $\star$ -moment limit of random matrices

$$(12) \quad A_N = D_N + \sqrt{o} T_N,$$

where D_N is a diagonal random matrix with eigenvalues $\lambda_1, \dots, \lambda_n$ which are i.i.d. random variables with distribution given by ν and

$$(13) \quad T_N = \begin{bmatrix} 0 & g_{1,2} & \cdots & g_{1,n-1} & g_{1,n} \\ 0 & 0 & \cdots & g_{2,n-1} & g_{2,n} \\ \vdots & & \ddots & \vdots & \vdots \\ 0 & & \cdots & 0 & g_{n-1,n} \\ 0 & & \cdots & 0 & 0 \end{bmatrix},$$

is an upper-triangular random matrix where $(\Re g_{i,j}, \Im g_{i,j})_{1 \leq i < j \leq N}$ are i.i.d. $N(0, \frac{1}{N})$ random variables. We recall that the Brown measure and offdiagonality of x are given by $\mu_x = \nu$ and $\text{od}_x = o$.

Theorem 9. *For any compactly supported probability measure ν on $\mathbb{C}$ and any number $o > 0$ if x is a $DT(\nu, o)$ then the inequality (11) becomes equality.*

Proof. Let us fix $R > 0$. Similarly as in Lemma 1 let (θ_N) be a sequence of non-negative numbers and (l_N) be a sequence of natural numbers such that

$$\lim_{N \rightarrow \infty} \theta_N = 0, \quad \lim_{N \rightarrow \infty} l_N = \infty,$$

$$\hat{\chi}_R^d(x) \leq \limsup_{N \rightarrow \infty} \hat{\chi}_R^d(x; N, l_N, \theta_N).$$

In definition (12) of A_N let us change D_N to be any (non-random) element of the set $\hat{\Gamma}_R^d(x; N, l_N, \theta_N)$. From results of Dykema and Haagerup [DH00] it follows that despite this change the sequence A_N still converges in expected $\star$ -moments to x :

$$(14) \quad \lim_{N \rightarrow \infty} \mathbb{E} \text{tr}(A_N^{s_1} \cdots A_N^{s_k}) = \phi(x^{s_1} \cdots x^{s_k})$$

for any $k \in \mathbb{N}$ and $s_1, \dots, s_k \in \{1, \star\}$ and by using similar combinatorial arguments as in [Tho00] one can show that

$$(15) \quad \lim_{N \rightarrow \infty} \text{Var} \text{tr}(A_N^{s_1} \cdots A_N^{s_k}) = 0.$$

Since $\limsup_{N \rightarrow \infty} \|T_N\| < \infty$ almost surely [Gem80], therefore there exists $R' > 0$ such that for any integer k and $\epsilon > 0$

$$(16) \quad \lim_{N \rightarrow \infty} P(\omega \in \Omega : D_N + \sqrt{o}T_N(\omega) \in \Gamma_{R'}(x; k, N, \epsilon)) = 1.$$

Furthermore since the convergence in (14) and (15) is uniform with respect to choice of the sequence (D_N) , hence it is possible to find universal R' for all choices of (D_N) .

By comparing the densities of two measures on $\mathcal{M}_N^{\text{sut}}$: the Lebesgue measure vol^{sut} and the distribution of the Gaussian random matrix $\sqrt{o}T_N$ we see that (16) implies that for every $0 < \delta < 1$, every integer k and $\epsilon > 0$ there exists N_0 such that for $N > N_0$ we have that the volume of the set

$\{m \in \mathcal{M}_N^{\text{sut}} : D_N + m \in \Gamma_{R'}(x; k, N, \epsilon)\}$ is bigger or equal to the volume of a $N(N-1)$ -dimensional ball with radius $\sqrt{(1-\delta)oN}$:

$$(17) \quad \text{vol}^{\text{sut}}\{m \in \mathcal{M}_N^{\text{sut}} : D_N + m \in \Gamma_{R'}(x; k, N, \epsilon)\} \geq \pi^{\frac{N(N-1)}{2}} \left[\Gamma\left(\frac{N(N-1)}{2} + 1\right) \right]^{-1} [(1-\delta)oN]^{\frac{N(N-1)}{2}}.$$

Since the sequence (D_N) was chosen arbitrarily, it follows that for every $0 < \delta < 1$, every integer k and $\epsilon > 0$ there exists N_0 such that for any $N > N_0$ and any $D_N \in \hat{\Gamma}_R^d(x; N, l_N, \theta_N)$ inequality (17) holds.

Now it is enough to apply Proposition 5 to show that for every $0 < \delta < 1$, every integer k and $\epsilon > 0$ we have

$$\chi_{R'}(x; k, \epsilon) \geq \hat{\chi}_R^d(x) + \frac{1}{2} + \frac{\log 2\pi(1-\delta)o}{2}$$

what finishes the proof. $\square$

4.6. Comparison of inequalities for free entropy. The inequality (11) contains a double integral, which is often called logarithmic energy of a measure. A similar term appears in the formula for the free entropy of a single selfadjoint operator which is due to Voiculescu [Voi94]:

$$(18) \quad \chi^{\text{sa}}(x) = \int_{\mathbb{R}} \int_{\mathbb{R}} \log |z_1 - z_2| d\mu_x(z_1) d\mu_x(z_2) + \frac{3}{4} + \frac{\log 2\pi}{2}.$$

It should be stressed that—despite the formal resemblance—the free entropies in (11) and (18) are different objects. Namely, in (11) we consider a free entropy $\chi(x)$ of a non-selfadjoint random variable while in (18) we consider a free entropy $\chi^{\text{sa}}(x)$ of a selfadjoint random variable which is defined by hermitian matrix approximations.

On the other hand inequality (11) contains a term equal to the logarithm of the offdiagonality, which can be regarded as a non-commutative variance. A similar expression appears in the Voiculescu's inequality for a free entropy of a non-selfadjoint variable [Voi94]:

$$(19) \quad \chi(x) \leq \log \left[\pi e (\phi(|x|^2) - |\phi(x)|^2)^2 \right].$$

5. ACKNOWLEDGEMENTS

The research was conducted at Texas A&M University on a scholarship funded by Polish-US Fulbright Commission. I acknowledge the support of Polish Research Committee grant No. P03A05415.

REFERENCES

- [Bro86] L. G. Brown. Lidskii's theorem in the type II case. In *Geometric methods in operator algebras (Kyoto, 1983)*, pages 1–35. Longman Sci. Tech., Harlow, 1986.
- [DH00] Kenneth Dykema and Uffe Haagerup. Invariant subspaces of Voiculescu's circular operator. *preprint*, 2000.
- [DH01] Kenneth Dykema and Uffe Haagerup. Decomposability of Voiculescu's circular operator and DT -operators. *preprint*, 2001.
- [FK52] Bent Fuglede and Richard V. Kadison. Determinant theory in finite factors. *Ann. of Math. (2)*, 55:520–530, 1952.
- [Ge97] Liming Ge. Applications of free entropy to finite von Neumann algebras. *Amer. J. Math.*, 119(2):467–485, 1997.
- [Gem80] Stuart Geman. A limit theorem for the norm of random matrices. *Ann. Probab.*, 8(2):252–261, 1980.
- [Haa01] Uffe Haagerup. Spectral decomposition of all operators in a II_1 factor, which is embedable in R^ω . *preprint*, 2001.
- [Had98] Don Hadwin. Free entropy and approximate equivalence in von Neumann algebras. In *Operator algebras and operator theory (Shanghai, 1997)*, pages 111–131. Amer. Math. Soc., Providence, RI, 1998.
- [Meh91] Madan Lal Mehta. *Random matrices*. Academic Press Inc., Boston, MA, second edition, 1991.
- [NSS99] Alexandru Nica, Dimitri Shlyakhtenko, and Roland Speicher. Maximality of the microstates free entropy for R -diagonal elements. *Pacific J. Math.*, 187(2):333–347, 1999.
- [Śni01] Piotr Śniady. Random regularization of Brown spectral measure. *Preprint. Available at <http://arXiv.org/abs/math/0105109>*, 2001.
- [Tho00] S. Thorbjørnsen. Mixed moments of Voiculescu's Gaussian random matrices. *J. Funct. Anal.*, 176(2):213–246, 2000.
- [Voi94] Dan Voiculescu. The analogues of entropy and of Fisher's information measure in free probability theory. II. *Invent. Math.*, 118(3):411–440, 1994.
- [Voi96] D. Voiculescu. The analogues of entropy and of Fisher's information measure in free probability theory. III. The absence of Cartan subalgebras. *Geom. Funct. Anal.*, 6(1):172–199, 1996.

INSTITUTE OF MATHEMATICS, UNIVERSITY OF WROCLAW, PL. GRUNWALDZKI
2/4, 50-384 WROCLAW, POLAND

E-mail address: `Piotr.Sniady@math.uni.wroc.pl`